

BLMH Rabbit pAb

货号: **AYP13425**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:20 - 1:50
理论分子量	52kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NCI-H460,Rat thymus
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human BLMH (NP_000377.1).
序列	MSSSGLNSEKVAALIQKLNSDPQFVLAQNVGTTDDLDDICLKRAVQRAQHVQHAVPQEGKIPITNQSSGRCWIFSLNVMRLPFMKKLNIEEFESQSYLFFWDKVERCYFFLSAFVDTAQRKEPEDGRLVQFLMNPANDGGQWDMLVNIVEKYGVI PKKCFPESYTTEATRRMNDILNHKMREFCIRLRNLVHSGATKGEISATQDVMMEEIFRVVCICLGNPPETFTWEYRDKDKNY QKIGPITPLEFYREHVKPLFNMEDKICLVNDPRPQHKYNKLYTVEYLSNMVGGKRTLY

靶点信息

研究背景	Bleomycin hydrolase (BMH) is a cytoplasmic cysteine peptidase that is highly conserved through evolution; however, the only known activity of the enzyme is metabolic inactivation of the glycopeptide bleomycin (BLM), an essential component of combination chemotherapy regimens for cancer. The protein contains the signature active site residues of the cysteine protease papain superfamily.
基因ID	642
基因名	BLMH
Swiss	Q13867
别名	BLMH;BH;BMH

产品验证



Western blot analysis of BLMH expressed in NCI-H460, Rat thymus using BLMH Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn