

# TNFRSF10A Rabbit pAb

货号: **AYP13457**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Homo sapiens
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	50kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	DU145
细胞定位	Membrane, Single-pass type I membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 269-468 of human TNFRSF10A (NP_003835.3).
序列	GGDPKCMDRVCFWRLGLLRGPGEADNAHNEILSNADSLSTFVSEQMESQEPADLTGVTVQSPGEAQCLLGPAEAEGS QRRRLVLPANGADPTETLMLFFDKFANIVPFDSWDQLMRQLDLTKNEIDVVRAGTAGPGDALYAMLKVVNKTGRNASI HTLLDALERMEERHAREKIQDLLVDSGKFIYLEDGTGSAVSLE

## 靶点信息

研究背景	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL), and thus transduces cell death signal and induces cell apoptosis. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein.
基因ID	8797

基因名	TNFRSF10A
Swiss	O00220
别名	TNFRSF10A;APO2;CD261;DR4;TRAILR-1;TRAILR1

## 产品验证



Western blot analysis of TNFRSF10A expressed in DU145 using TNFRSF10A Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)