

RICTOR Rabbit pAb

货号: **AYP13467**

产品信息

反应	Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens, Mus musculus
应用	WB
推荐浓度	WB: 1:100 - 1:500
理论分子量	28kDa/192kDa/195kDa
实测分子量	200KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	PC-12
细胞定位	cytosol
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1290-1500 of human RICTOR (NP_689969.2).
序列	PGSSHTLPRAQSLKAPSIATIKSLADCNFSYTSSRDAFGYATLKRLLQQRMHPSLSHSEALASPAKDVLFDTITMKANSF ESRLTPSRFMKALSYASLDKEDLLSPINQNTLQRSSSVRSMVSSATYGGSDDYIGLALPVDINDIFQVKDIPYFQTKNIPPH DDRGARAFAHADAGGLPSGTGGLVKNSFHLLRQQMSLTEIMNSIHSDA

靶点信息

研究背景	RICTOR and MTOR (FRAP1; MIM 601231) are components of a protein complex that integrates nutrient- and growth factor-derived signals to regulate cell growth (Sarbasov et al., 2004 [PubMed 15268862]). [supplied by OMIM, Mar 2008]
基因ID	253260

基因名	RICTOR
Swiss	Q6R327
别名	RICTOR;AVO3;PIA;hAVO3

产品验证



Western blot analysis of RICTOR expressed in PC-12 using RICTOR Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn