

— ABLYBIO, Help Your Research



NOGO Receptor 1 (RTN4R) Rabbit pAb

货号: AYP13486

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens IHC: Homo sapiens
应用	WB IF/ICC
推荐浓度	WB: 1:100 - 1:500 IF/ICC: 1:50 - 1:200
理论分子量	50kDa
实测分子量	66KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	C6,Rat uterus
细胞定位	Cell membrane,GPI-anchor,Lipid-anchor
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 338-410 of human NOGO Receptor 1 (RTN4R) (NP_075380.1).
------	---

靶点信息

研究背景	This gene encodes the receptor for reticulon 4, oligodendrocyte myelin glycoprotein and myelin-associated glycoprotein. This receptor mediates axonal growth inhibition and may play a role in regulating axonal regeneration and plasticity in the adult central nervous system.
基因ID	65078
基因名	RTN4R
Swiss	Q9BZR6 (https://www.uniprot.org/uniprotkb/Q9BZR6/entry)
别名	RTN4R,NGR,NOGOR,NOGO Receptor 1 (RTN4R) Rabbit pAb,Nogo receptor,Nogo-66 receptor

产品验证



Western blot analysis of NOGO Receptor 1 (RTN4R) expressed in C6, Rat uterus using NOGO Receptor 1 (RTN4R) Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)