

# FGG Rabbit pAb

货号: **AYP13516**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Mus musculus
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	49kDa/51kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver
细胞定位	Secreted
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 50-330 of human FGG (NP_000500.2).
序列	GIADFLSTYQTKVDKDLQSLEDILHQVENKTSEVKQLIKAIQLTYNPDESSKPNMIDAATLKS RKMLEEIMKYEASILTHDSSI RYLQEIYNSNNQKIVNLKEKVAQLEAQCQEPCKDVTQIHDITGKDCQDIANKGAKQSGLYFIKPLKANQQFLVYCEIDGSG NGWTVFQKRLDGSVDFKKNWIQYKEGFGHLSPTGTTEFWLGNKIHILISTQSAIPYALRVELEDWNGRTSTADYAMFKV GPEADKYRLTYAYFAGGDAGDAFDGDFGDDPSDKFF

## 靶点信息

研究背景	The protein encoded by this gene is the gamma component of fibrinogen, a blood-borne glycoprotein comprised of three pairs of nonidentical polypeptide chains. Following vascular injury, fibrinogen is cleaved by thrombin to form fibrin which is the most abundant component of blood clots. In addition, various cleavage products of fibrinogen and fibrin regulate cell adhesion and spreading, display vasoconstrictor and chemotactic activities, and are mitogens for several cell types. Mutations in this gene lead to several disorders, including dysfibrinogenemia, hypofibrinogenemia and thrombophilia. Alternative splicing results in transcript variants encoding different isoforms.
基因ID	2266
基因名	FGG
Swiss	P02679
别名	FGG

## 产品验证



Western blot analysis of FGG expressed in Mouse liver using FGG Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)