

# BPI Rabbit pAb

货号: **AYP13559**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>IF:</b> Homo sapiens,Mus musculus
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	53kDa
实测分子量	55kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,HepG2,SK-Hep1,Mouse liver,Mouse thymus
细胞定位	Cytoplasmic granule membrane,Secreted
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 248-487 of human BPI ( NP_001716.2).
序列	AETLDVQMKGEFYSENHHNPPPFAPPVMEFPAAHDRMVYLGSLDYFFNTAGLVYQEAGVLKMTLRDDMIPKESKFRLLTK FFGTFLPEVAKKFPNMKIQIHVSASTPPHLSVQPTGLTFYPAVDVQAFVLPNSSLASLFLIGMHTTGSMESVAESNRLVGE LKLDRLLLLKHSNIGPPFVELLQDIMNYIVPILVLPVNEKLQKGFPLTPARVQLYNVVLQPHQNFLFLGADVVK

## 靶点信息

研究背景	This gene encodes a lipopolysaccharide binding protein. It is associated with human neutrophil granules and has antimicrobial activity against gram-negative organisms.
基因ID	671

基因名	BPI
Swiss	P17213
别名	BPI;BPIFD1;rBPI

## 产品验证



Western blot analysis of BPI expressed in Mouse liver, Jurkat, HepG2 using BPI Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)