

# BDH1 Rabbit pAb

货号: **AYP13611**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Mus musculus
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:1000 - 1:2000
理论分子量	38kDa
实测分子量	42kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,MCF7,SKOV3,LO2
细胞定位	Mitochondrion matrix
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-343 of human BDH1 ( NP_004042.1).
序列	MLATRLSRPLSRLPGKTLACDRENGARRPLLLGSTSFIPIGRRTYASAAEPVGSKAVLVTGCDSGFGFSLAKHLHSGFLV FAGCLMKDKGHDGVKELDSLNSDRLRTVQLNVCSSSEVEKVVIVRSSLKDPEKGMWGLVNNAGISTFGEVEFTSLETYK QVAEVLWGTVRMTKSFLLPILIRRAKGRVVNISSMLGRMANPARSPYCITKFGVEAFSDCLRYEMYPLGVKVSVEPGNFIA ATSLYSPESIQAIKKMWEELPEVVRKDYGKKYFDEKIAKMETYCSSGSTDTSPVIDAVTHALTATTPYTRYHPMDYYWWL RMQIMTHLPGAISDMIYIR

## 靶点信息

研究背景	This gene encodes a member of the short-chain dehydrogenase/reductase gene family. The encoded protein forms a homotetrameric lipid-requiring enzyme of the mitochondrial membrane and has a specific requirement for phosphatidylcholine for optimal enzymatic activity. The encoded protein catalyzes the interconversion of acetoacetate and (R)-3-hydroxybutyrate, the two major ketone bodies produced during fatty acid catabolism. Alternatively spliced transcript variants encoding the same protein have been described.
基因ID	622
基因名	BDH1
Swiss	Q02338
别名	BDH1;BDH;SDR9C1

## 产品验证



Western blot analysis of BDH1 expressed in A-549, MCF7, SKOV3 using BDH1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)