

CDKN2AIP Rabbit pAb

货号: **AYP13628**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:100
理论分子量	61kDa
实测分子量	65kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SKOV3,Mouse thymus,Rat thymus
细胞定位	Nucleus,nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-220 of human CDKN2 AIP (NP_060102.1).
序列	MAQEVSEYLSQNPRVAAWVEALRCDDGETDKHWRHRRDFLLRNAGDLAPAGGAASASTDEAADAESGTRNRQLQLISF SMAWANHVFLGCRYPQKVMKDILSMAEGIKVTDAPTYTTRDELVAKVKKRGISSSNEGVEEPSKKRVIEGKNSSAVEQDH AKTSAKTERASAAQQENSSTCIGSAIKSESGNSARSSGISSQNSSTSDGDRSVSSQSSSSVSS

靶点信息

研究背景	The protein encoded by this gene regulates the DNA damage response through several different signaling pathways. One such pathway is the p53-HDM2-p21(WAF1) pathway, which is critical to the DNA damage response. Two transcript variants encoding different isoforms have been found for this gene.
基因ID	55602

基因名	CDKN2AIP
Swiss	Q9NXV6
别名	CDKN2AIP;CARF

产品验证



Western blot analysis of CDKN2AIP expressed in SKOV3, Mouse thymus, Rat thymus using CDKN2AIP Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn