

— ABLYBIO, Help Your Research



SARS-CoV-2 NSP9 Rabbit pAb

货号: **AYP13645**

产品信息

| | |
|-------|---|
| 反应 | SARS-CoV-2 |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | IF: Homo sapiens IP: Homo sapiens |
| 应用 | WB |
| 推荐浓度 | WB: 1:2000 - 1:6000 |
| 理论分子量 | |
| 实测分子量 | 15-20KDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | 293T |
| 细胞定位 | [Host translation inhibitor nsp1]: Host cytoplasm, [Non-structural protein 2]: Host cytoplasm, [Papain-like protease nsp3]: Host endoplasmic, [Non-structural protein 4]: Host endoplasmic, [3C-like proteinase nsp 5]: Host cytoplasm, [Non-structural protein 6]: Host endoplasmic, [Non-structural protein 7]: Host cytoplasm, host, [Non-structural protein 8]: Host cytoplasm, [Viral protein genome-linked nsp9]: Host, [Non-structural protein 10]: Host cytoplasm, [Guanine-N7 methyltransferase nsp14]: Host, [Helicase nsp13]: Host endoplasmic reticulum-, [Uridylate-specific endoribonuclease nsp15]: Host, [2'-O-methyltransferase nsp16]: Host nucleus |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-113 of coronavirus NSP9 (YP_009742616.1). |
|------|---|

靶点信息

| | |
|-------|--|
| 研究背景 | [Replicase polyprotein 1ab]: Multifunctional protein involved in the transcription and replication of viral RNAs. Contains the proteinases responsible for the cleavages of the polyprotein. |
| 基因ID | 43740578 |
| 基因名 | rep |
| Swiss | P0DTD1 (https://www.uniprot.org/uniprotkb/P0DTD1/entry) |
| 别名 | SARS-CoV-2 NSP9 Rabbit pAb,rep,ORF1ab polyprotein,Leader protein,Non-structural protein 1,p65 homolog,Non-structural protein 3,PL2-PRO,Papain-like proteinase,Main protease,Non-structural protein 5,SARS coronavirus main proteinase,Non-structural protein 9 |

产品验证



Western blot analysis of SARS-CoV-2 NSP9 expressed in 293T using SARS-CoV-2 NSP9 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)