

IMMT Rabbit pAb

货号: **AYP13667**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens,Mus musculus IP: Homo sapiens,Mus musculus
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	80kDa/82kDa/83kDa
实测分子量	90KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,293T,A-549,Mouse brain
细胞定位	Mitochondrion,Mitochondrion inner membrane,Single-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 120-270 of human IMMT (NP_006830.2).
序列	ESKQPASQLQKQKGDTPASATAPTEAAQIISAAGDTLSVPAPAVQPEESLKTDPHEIGEGKPTPALSEEASSSSIRERPPEEV AARLAQQEQEKQEVKIESLAKSLEDALRQTASVTLQAIQAQNAAVQAVNAHSNILKAAMDNSEIAGEKK

靶点信息

研究背景	Component of the MICOS complex, a large protein complex of the mitochondrial inner membrane that plays crucial roles in the maintenance of crista junctions, inner membrane architecture, and formation of contact sites to the outer membrane. Plays an important role in the maintenance of the MICOS complex stability and the mitochondrial cristae morphology.
------	---

基因ID	10989
基因名	IMMT
Swiss	Q16891
别名	IMMT;HMP;MINOS2;Mic60;P87;P87/89;P89;PIG4;PIG52

产品验证



Western blot analysis of IMMT expressed in Mouse brain,HT-29,293T using IMMT Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn