

# HSF2 Rabbit pAb

货号: **AYP13668**

## 产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Homo sapiens
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	58kDa/60kDa
实测分子量	75KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3,RAW264.7,Rat testis
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 39-215 of human HSF2 ( Q03933).
序列	LDEQRFAKEILPKYFKHNNMASFVRQLNMYGFRKVVHIDSGIVKQERDGPVEFQHPYFKQGDDLLENIKRKVSSSKPEE NKIRQEDLTKIISSAQKVQIKQETIESRLESELKSENE <sup>SLWKEVSELRAKHAQQQVIRKIVQFIVTLVQNNQLVSLKRRKRP</sup> LLL NTNGAQKKNL <sup>FQH</sup>

## 靶点信息

研究背景	The protein encoded by this gene belongs to the HSF family of transcription factors that bind specifically to the heat-shock promoter element and activate transcription. Heat shock transcription factors activate heat-shock response genes under conditions of heat or other stresses. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
基因ID	3298

基因名	HSF2
Swiss	Q03933
别名	HSF2;HSF 2;HSTF 2;HSTF2

## 产品验证



Western blot analysis of HSF2 expressed in NIH/3T3,RAW264.7,Rat testis using HSF2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)