

# DKK1 Rabbit pAb

货号: **AYP13680**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Homo sapiens
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	28kDa
实测分子量	28-45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,K-562,Mouse brain,Rat brain,Rat liver
细胞定位	Secreted
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 32-266 of human DKK1 (NP_036374.1).
序列	TLNSVLNSNAIKNLPPLGGAAGHPGSAVSAAPGILYPGGNKYQTIDNYQPYPCAEDECCTDEYCASPTRGGDAGVQICL ACRKRKRRCMRHAMCCPGNYCKNGICVSSDQNHFRGEIETITESFGNDHSTLDGYSRRTTLSSKMYHTKGQEGSVCLR SSDCASGLCCARHFWSKICKPVLKEGVCTKHRRKGGSHGLEIFQRCYCGEGLSRIQKDHQASNSSRLHTCQRH

## 靶点信息

研究背景	This gene encodes a protein that is a member of the dickkopf family. It is a secreted protein with two cysteine rich regions and is involved in embryonic development through its inhibition of the WNT signaling pathway. Elevated levels of DKK1 in bone marrow plasma and peripheral blood is associated with the presence of osteolytic bone lesions in patients with multiple myeloma.
基因ID	22943

基因名	DKK1
Swiss	O94907
别名	DKK1;DKK-1;SK

## 产品验证



Western blot analysis of DKK1 expressed in Mouse brain,Rat brain,HeLa using DKK1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)