

# Uhrf2 Rabbit pAb

货号: **AYP13695**

## 产品信息

|       |  |
|-------|--|
| 反应    | Human,Mouse,Rat  |
| 宿主    | Rabbit   |
| 克隆性   | Polyclonal   |
| 预测反应  | <b>WB:</b> Homo sapiens  |
| 应用    | <a href="#">WB</a>   |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:1000  |
| 理论分子量 | 56kDa/89kDa  |
| 实测分子量 | 95KDa  |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated   |
| 阳性对照  | Mouse thymus   |
| 细胞定位  | Nucleus  |
| 纯化    | Affinity purification  |

## 抗原信息


|      |   |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 636-733 of mouse Uhrf2 (NP_659122.2). |
| 序列   | LQYPAGYPSEKEGKKTGQSKKQGSEATKRPASDDECPGDSKVLKASDSTDAVEAFQLTPQQRLIREDCQNQKLWDE<br>VLASLVEGPNFLKKLEQSFM                |

## 靶点信息

|      |   |
|------|---|
| 研究背景 | This gene encodes a nuclear protein which is involved in cell-cycle regulation. The encoded protein is a ubiquitin-ligase capable of ubiquinating PCNP (PEST-containing nuclear protein), and together they may play a role in tumorigenesis. The encoded protein contains an NIRF_N domain, a PHD finger, a set- and ring-associated (SRA) domain, and a RING finger domain and several of these domains have been shown to be essential for the regulation of cell proliferation. This protein may also have a role in intranuclear degradation of polyglutamine aggregates. Alternative splicing results in multiple transcript variants some of which are non-protein coding. |
|------|---|

|       |                                     |
|-------|-------------------------------------|
| 基因ID  | 109113                              |
| 基因名   |                                     |
| Swiss | Q7TMI3                              |
| 别名    | UHRF2;NIRF;RNF107;TDRD23;URF2;Uhrf2 |

## 产品验证



Western blot analysis of Uhrf2 expressed in Mouse thymus using Uhrf2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)