

DLGAP5 Rabbit pAb

货号: **AYP13707**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	85kDa/94kDa/95kDa
实测分子量	120kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562
细胞定位	Cytoplasm,Nucleus,cytoskeleton,spindle
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 547-846 of human DLGAP5 (NP_055565.3).
序列	KNVFRKKVVGSIASKPKQDDAGRIAARNRLAAIKNAMRERIRQECAETAVSVIPKEVDKIVFDAGFFRVESPVKLFSGLSV SSEGPSQRLGTPKSVNKAVSQSRNEMGIPQQTTSPEAGPQNTKSEHVKKTLFLSIPESRSSIEDAQCPGLPDLIEENHVV NKTDLKVDCLESSERMSLPLLAGGVADDINTNKKEGISDVVEGMELNSSITSQDVLMSSEKNTASQNSILEEGETKISQSE LFDNKSLTTECHLLDSPGLNCSNPFTQLERRHQEHARHISFGGNLITFSPLQPGEF

靶点信息

研究背景	Potential cell cycle regulator that may play a role in carcinogenesis of cancer cells. Mitotic phosphoprotein regulated by the ubiquitin-proteasome pathway. Key regulator of adherens junction integrity and differentiation that may be involved in CDH1-mediated adhesion and signaling in epithelial cells.
------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

基因ID	9787
基因名	DLGAP5
Swiss	Q15398
别名	DLGAP5;DLG7;HURP

产品验证



Western blot analysis of DLGAP5 expressed in K-562 using DLGAP5 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn