

# IFI35 Rabbit pAb

货号: **AYP13729**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Mus musculus
应用	<a href="#">WB</a> <a href="#">IF/ICC</a>
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:100
理论分子量	31kDa
实测分子量	35kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HepG2,A-431
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-288 of human IFI35 (NP_005524.2).
序列	MSAPLDAALHALQEEQARLKMRLWDLQQLRKELGDSPKDKVPFVSPKIPLVFRGHTQQDPEVPKSLVSNLRIHCPLLAGS ALITFDDPKVAEQVLQQKEHTINMEECRLRVQVQPLELPMVTTIQVMMSSQLSGRRVLVTGFPAASLRLSEEELDKLEIFFG KTRNGGGDVDVRELLPGSVMLGFARDGVAQRLCQIGQFTVPLGGQQVPLRVSPYVNGEIQKAEIRSQPVPRSVLVLNIPDI LDGPELHDVLEIHFQKPTRGGGEVEALTVVPQGQQGLAVFTSESG

## 靶点信息

研究背景	Acts as a signaling pathway regulator involved in innate immune system response. In response to interferon IFN-alpha, associates in a complex with signaling pathway regulator NMI to regulate immune response ; the complex formation prevents proteasome-mediated degradation of IFI35 and correlates with IFI35 dephosphorylation. In complex with NMI, inhibits virus-triggered type I interferon/IFN-beta production. In complex with NMI, negatively regulates nuclear factor NF-kappa-B signaling by inhibiting the nuclear translocation, activation and transcription of the NF-kappa-B subunit p65/RELA, resulting in the inhibition of endothelial cell proliferation, migration and re-endothelialization of injured arteries. Beside its role as an intracellular signaling pathway regulator, also functions extracellularly as damage-associated molecular patterns (DAMPs) to promote inflammation when actively released by macrophage to the extracellular space during cell injury and pathogen invasion. Macrophage-secreted IFI35 activates NF-kappa-B signaling in adjacent macrophages through Toll-like receptor 4/TLR4 activation, thereby inducing NF-kappa-B translocation from the cytoplasm into the nucleus which promotes the release of proinflammatory cytokines.
基因ID	3430
基因名	IFI35
Swiss	P80217
别名	IFI35;IFP35

## 产品验证



Western blot analysis of IFI35 expressed in HeLa, HepG2, A-431 using IFI35 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)