

TMEM100 Rabbit pAb

货号: **AYP13748**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens IF: Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	14kDa
实测分子量	14kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,SH-SY5Y,Mouse lung
细胞定位	endoplasmic reticulum,perikaryon,perinuclear region of cytoplasm,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 50 to the C-terminus of human TME M100 (NP_001093110.1).
序列	ELSCYRCIIPFAVVVFIAGIVVTAVAYSFNSHGSIIIFGLVVLSSGLFLLASSALCWKVRQRSKKAKRRESQ TALVANQRS LFA

靶点信息

研究背景	Plays a role during embryonic arterial endothelium differentiation and vascular morphogenesis through the ACVRL1 receptor-dependent signaling pathway upon stimulation by bone morphogenetic proteins, such as GDF2/BMP9 and BMP10. Involved in the regulation of nociception, acting as a modulator of the interaction between TRPA1 and TRPV1, two molecular sensors and mediators of pain signals in dorsal root ganglia (DRG) neurons. Mechanistically, it weakens their interaction, thereby releasing the inhibition of TRPA1 by TRPV1 and increasing the single-channel open probability of the TRPA1-TRPV1 complex.
------	---

基因ID	55273
基因名	TMEM100
Swiss	Q9NV29
别名	TMEM100

产品验证



Western blot analysis of TMEM100 expressed in 293T,SH-SY5Y,Mouse lung using TMEM100 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn