

KCNN4 Rabbit pAb

货号: **AYP13767**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	IHC: Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	47kDa
实测分子量	53kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HepG2
细胞定位	Cell membrane, Multi-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 288-427 of human KCNN4 (NP_002241.1).
序列	KLEFNKAEKHVHNFMMDIQYTKEMKESAAARVLQEAWMFYKHTRRKESHAARRHQRKLLAAINAFRQVRLKHRKLREQVN SMVDISKMHMILYDLQNLSSSHRALEKQIDTLAGKLDALTELLSTALGPRQLPEPSQSK

靶点信息

研究背景	The protein encoded by this gene is part of a potentially heterotetrameric voltage-independent potassium channel that is activated by intracellular calcium. Activation is followed by membrane hyperpolarization, which promotes calcium influx. The encoded protein may be part of the predominant calcium-activated potassium channel in T-lymphocytes. This gene is similar to other KCNN family potassium channel genes, but it differs enough to possibly be considered as part of a new subfamily.
基因ID	3783

基因名	KCNN4
Swiss	O15554
别名	KCNN4;DHS2;IK;IK1;IKCA1;KCA4;KCa3.1;SK4;hIKCa1;hKCa4;hSK4

产品验证



Western blot analysis of KCNN4 expressed in HepG2 using KCNN4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn