

— ABLYBIO, Help Your Research



SMARCC2 Rabbit pAb

货号: **AYP13771**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB
推荐浓度	WB: 1:200 - 1:2000
理论分子量	124kDa/126kDa/132kDa
实测分子量	171kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,Mouse spinal cord
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 150-250 of human SMARCC2 (NP_620706.1).
------	--

靶点信息

研究背景	The protein encoded by this gene is a member of the SWI/SNF family of proteins, whose members display helicase and ATPase activities and which are thought to regulate transcription of certain genes by altering the chromatin structure around those genes. The encoded protein is part of the large ATP-dependent chromatin remodeling complex SNF/SWI and contains a predicted leucine zipper motif typical of many transcription factors. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
基因ID	6601
基因名	SMARCC2
Swiss	Q8TAQ2 (https://www.uniprot.org/uniprotkb/Q8TAQ2/entry)
别名	SMARCC2,BAF170,CRACC2,Rsc8,SMARCC2 Rabbit pAb,BRG1-associated factor 170,SWI/SNF complex 170 kDa subunit,SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily C member 2

产品验证



Western blot analysis of SMARCC2 expressed in U-87MG, Mouse spinal cord using SMARCC2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)