

SMARCC2 Rabbit pAb

货号: **AYP13771**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | WB: Mus musculus |
| 应用 | WB |
| 推荐浓度 | WB: 1:200 - 1:2000 |
| 理论分子量 | 124kDa/126kDa/132kDa |
| 实测分子量 | 171kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | U-87MG,Mouse spinal cord |
| 细胞定位 | Nucleus |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | A synthetic peptide corresponding to a sequence within amino acids 150-250 of human SMARCC2 (NP_620706.1). |
| 序列 | PKLLGKLDIIKRHQGTVTEDKNNASHVVYPVPGNLEEEWVRPVMKRDQVLLHWGYYPDSYDTWIPASEIEASVEDAP TPEKPRKVHAKWILDTFNE |

靶点信息

| | |
|------|---|
| 研究背景 | The protein encoded by this gene is a member of the SWI/SNF family of proteins, whose members display helicase and ATPase activities and which are thought to regulate transcription of certain genes by altering the chromatin structure around those genes. The encoded protein is part of the large ATP-dependent chromatin remodeling complex SNF/SWI and contains a predicted leucine zipper motif typical of many transcription factors. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. |
|------|---|

| | |
|-------|----------------------------|
| 基因ID | 6601 |
| 基因名 | SMARCC2 |
| Swiss | Q8TAQ2 |
| 别名 | SMARCC2;BAF170;CRACC2;Rsc8 |

产品验证



Western blot analysis of SMARCC2 expressed in U-87MG, Mouse spinal cord using SMARCC2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn