

CXCL9 Rabbit pAb

货号: **AYP13779**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	14kDa
实测分子量	14kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	THP-1, A-431, 293T
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 50 to the C-terminus of human CXCL9 (NP_002407.1).
序列	KQFAPSPSCEKIEIATLKNGVQTCLNPDSADVKELIKKWEKQVSQKKKQKNGKKHQKKVLKVRKSQRSRQKKT

靶点信息

研究背景	This antimicrobial gene encodes a protein thought to be involved in T cell trafficking. The encoded protein binds to C-X-C motif chemokine 3 and is a chemoattractant for lymphocytes but not for neutrophils.
基因ID	4283
基因名	CXCL9

Swiss	Q07325
别名	CXCL9;CMK;Humig;MIG;SCYB9;crg-10

产品验证



Western blot analysis of CXCL9 expressed in THP-1,A-431,293T using CXCL9 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn