

RGS4 Rabbit pAb

货号: **AYP13787**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:100 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	10kDa/21kDa/23kDa/33kDa
实测分子量	30kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,HeLa,A-375,Jurkat,Raji
细胞定位	cytoplasm,cytosol,nucleus,plasma membrane
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human RGS4 (NP_005604.1).
序列	MCKGLAGLPASCLRSKMDMKHRLGFLQKSDSCEHNSSHNNKDKVVICQVRSQEEVKKWAESLENLISHECGLAAFKAFLLKSEYSEENIDFWISCEEYKK

靶点信息

研究背景	Regulator of G protein signaling (RGS) family members are regulatory molecules that act as GTPase activating proteins (GAPs) for G alpha subunits of heterotrimeric G proteins. RGS proteins are able to deactivate G protein subunits of the Gi alpha, Go alpha and Gq alpha subtypes. They drive G proteins into their inactive GDP-bound forms. Regulator of G protein signaling 4 belongs to this family. All RGS proteins share a conserved 120-amino acid sequence termed the RGS domain. Regulator of G protein signaling 4 protein is 37% identical to RGS1 and 97% identical to rat Rgs4. This protein negatively regulate signaling upstream or at the level of the heterotrimeric G protein and is localized in the cytoplasm. Alternatively spliced transcript variants have been found for this gene.
基因ID	5999
基因名	RGS4
Swiss	P49798
别名	RGS4;RGP4;SCZD9

产品验证



Western blot analysis of RGS4 expressed in U-251MG, HeLa, A-375 using RGS4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn