

GTF2B Rabbit pAb

货号: **AYP13799**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	34kDa
实测分子量	32KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Mouse thymus,Rat testis
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 40-154 of GTF2B (NP_001505.1).
序列	VVGDRVIDVGSEWRFTSNDKATKDP SRVGD SQNPLLS DGD LST MIGKGTGAASFDEFGNSKYQNRRTMSSSDRAMMN AFKEITTMADRINLPRNIVDR TN NLFKQVYEQKSLKGR

靶点信息

研究背景	This gene encodes the general transcription factor IIB, one of the ubiquitous factors required for transcription initiation by RNA polymerase II. The protein localizes to the nucleus where it forms a complex (the DAB complex) with transcription factors IID and IIA. Transcription factor IIB serves as a bridge between IID, the factor which initially recognizes the promoter sequence, and RNA polymerase II.
------	--

基因ID	2959
基因名	GTF2B
Swiss	Q00403
别名	GTF2B;TF2B;TFIIB

产品验证



Western blot analysis of GTF2B expressed in HeLa, Mouse thymus, Rat testis using GTF2B Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn