— ABLYBIO, Help Your Research

ABCD2 Rabbit pAb

货号: AYP13829

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	**WB:** Mus musculus
应用	WB IF/ICC
推荐浓度	**WB:** 1:500 - 1:2000 **IF/ICC:** 1:50 - 1:100
理论分子量	83kDa
实测分子量	83kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	mouse cerebellum tissue
细胞定位	Multi-pass membrane protein,Peroxisome membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 420-500 of human ABCD2 (NP_005155.1).

靶点信息

研究背景	The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the ALD subfamily, which is involved in peroxisomal import of fatty acids and/or fatty acyl-CoAs in the organelle. All known peroxisomal ABC transporters are half transporters which require a partner half transporter molecule to form a functional homodimeric or heterodimeric transporter. The function of this peroxisomal membrane protein is unknown; however this protein is speculated to function as a dimerization partner of ABCD1 and/or other peroxisomal ABC transporters. Mutations in this gene have been observed in patients with adrenoleukodystrophy, a severe demyelinating disease. This gene has been identified as a candidate for a modifier gene, accounting for the extreme variation among adrenoleukodystrophy phenotypes. This gene is also a candidate for a complement group of Zellweger syndrome, a genetically heterogeneous disorder of peroxisomal biogenesis.
基因ID	225
基因名	ABCD2
Swiss	Q9UBJ2 (https://www.uniprot.org/uniprotkb/Q9UBJ2/entry)
别名	ABCD2,ABC39,ALDL1,ALDR,ALDRP,hALDR,ABCD2 Rabbit pAb,Adrenoleukodystrophy-like 1,Adrenoleukodystrophy-related protein,ALD1

产品验证



Western blot analysis of ABCD2 expressed in mouse cerebellum tissue using ABCD2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)