

# ZNRF3 Rabbit pAb

货号: **AYP13831**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Homo sapiens <b>IHC:</b> Homo sapiens
应用	<a href="#">WB</a> <a href="#">IHC</a>
推荐浓度	<b>WB:</b> 1:1000 - 1:5000 <b>IHC:</b> 1:50 - 1:200
理论分子量	100kDa
实测分子量	101KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse testis
细胞定位	plasma membrane
纯化	Affinity purification


## 抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 50-150 of human ZNRF3 (NP_001193927.1).
序列	PGAARAKETAFVEVVLFESSPSGDYTTYTTGLTGRFSRAGATLSAEGEIVQMHPGLGCNNNDEEDLYEYGWVGVVVKLEQP ELDPKPCLTVLGKAKRAVQRG

## 靶点信息

研究背景	E3 ubiquitin-protein ligase that acts as a negative regulator of the Wnt signaling pathway by mediating the ubiquitination and subsequent degradation of Wnt receptor complex components Frizzled and LRP6. Acts on both canonical and non-canonical Wnt signaling pathway. Acts as a tumor suppressor in the intestinal stem cell zone by inhibiting the Wnt signaling pathway, thereby restricting the size of the intestinal stem cell zone. Along with RSPO2 and RNF43, constitutes a master switch that governs limb specification (By similarity).
基因ID	84133
基因名	ZNRF3
Swiss	Q9ULT6
别名	ZNRF3;BK747E2.3;RNF203

## 产品验证



Western blot analysis of ZNRF3 expressed in Mouse testis using ZNRF3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)