

GPR109A/HM74A/HCAR2 Rabbit pAb

货号: **AYP13875**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Sus scrofa IF/ICC: Sus scrofa
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	42kDa
实测分子量	44KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BxPC-3,Rat liver
细胞定位	cell junction,plasma membrane
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 100-200 of human GPR109A/HM74A/HCAR2 (NP_808219.1).
序列	CRLMLFMLAMNRQGSIIFLTVAVDYFRVWHPHHLNKKISNRATAIISCLLWGITIGLTVHLLKKKMPIQNGGANLCSFSICHTFQWHEAMFLLEFFLP

靶点信息

研究背景	Acts as a high affinity receptor for both nicotinic acid (also known as niacin and (D-beta-hydroxybutyrate and mediates increased adiponectin secretion and decreased lipolysis through G(i)-protein-mediated inhibition of adenylyl cyclase. This pharmacological effect requires nicotinic acid doses that are much higher than those provided by a normal diet. Mediates nicotinic acid-induced apoptosis in mature neutrophils. Receptor activation by nicotinic acid results in reduced cAMP levels which may affect activity of cAMP-dependent protein kinase A and phosphorylation of target proteins, leading to neutrophil apoptosis. The rank order of potency for the displacement of nicotinic acid binding is 5-methyl pyrazole-3-carboxylic acid = pyridine-3-acetic acid > acifran > 5-methyl nicotinic acid = acipimox >> nicotinuric acid = nicotinamide.
基因ID	338442
基因名	HCAR2
Swiss	Q8TDS4
别名	HCAR2;GPR109A;HCA2;HM74a;HM74b;NIACR1;PUMAG;Puma-g

产品验证



Western blot analysis of GPR109A/HM74A/HCAR2 expressed in BxPC-3, Rat liver using GPR109A/HM74A/HCAR2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn