

ULK2 Rabbit pAb

货号: **AYP13900**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Trionyx Sinensis
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	112kDa
实测分子量	110KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SKOV3,Mouse kidney,Rat lung,Rat brain
细胞定位	Cytoplasmic vesicle membrane,Peripheral membrane protein
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human ULK2 (NP_00113608 2.1).
序列	MEVVGDFEYSKRDLVGHGAFVAVFRGRHRQKTDWEVAIKSINKKNLSKSQLLGKEIKILKELQHENIVALYDVQELPNSVF LVMEYCNGGDLADYLQAK

靶点信息

研究背景	This gene encodes a protein that is similar to a serine/threonine kinase in <i>C. elegans</i> which is involved in axonal elongation. The structure of this protein is similar to the <i>C. elegans</i> protein in that both proteins have an N-terminal kinase domain, a central proline/serine rich (PS) domain, and a C-terminal (C) domain. The gene is located within the Smith-Magenis syndrome region on chromosome 17. Alternatively spliced transcripts encoding the same protein have been identified.
基因ID	9706

基因名	ULK2
Swiss	Q8IYT8
别名	ULK2;ATG1B;Unc51.2

产品验证



Western blot analysis of ULK2 expressed in Mouse kidney,Rat lung,SKOV3 using ULK2 Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn