

JDP2 Rabbit pAb

货号: **AYP13905**

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	18kDa/19kDa
实测分子量	18kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,HepG2,Rat testis
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-174 of human JDP2 (NP_001128521.1).
序列	MVAGWPATPPAMMPGQIPDPSVTTGSLPGLGPLTGLPSSALTVEELKYADIRNLGAMIAPLHFLEVKLGKRPQPVKSELDEEEERRKRRREKNKVAARCRNKKKERTEFLQRESERLELMNAELKTQIEELKQERQQLILMLNRHRPTCIVRTDSVKTPESG NPLLEQLEKK

靶点信息

研究背景	Component of the AP-1 transcription factor that represses transactivation mediated by the Jun family of p roteins. Involved in a variety of transcriptional responses associated with AP-1 such as UV-induced apopt osis, cell differentiation, tumorigenesis and antitumogeneris. Can also function as a repressor by recruitin g histone deacetylase 3/HDAC3 to the promoter region of JUN. May control transcription via direct regulat ion of the modification of histones and the assembly of chromatin.
------	---

基因ID	122953
基因名	JDP2
Swiss	Q8WYK2
别名	JDP2;JUNDM2

产品验证



Western blot analysis of JDP2 expressed in Rat testis, HeLa, Jurkat using JDP2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn