

CHKA Rabbit pAb

货号: **AYP13917**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	50kDa/52kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,Jurkat,Mouse brain,Rat testis
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-170 of human CHKA (NP_001268.2).
序列	MKTKFCTGGEAEPSPLGLLLSCGSGSAAPAGVGQQRDAASDLESKQLGGQPPLALPPPPLPLPLPQPPPPQPPAD EQPEPRTRRRAYLWCKEFLPGAWRGLREDEFHISVIRGGLSNMLFQCSPDATTATLGDEPRKVLRLLYGAILQMRSCKE SEQAQKENE

靶点信息

研究背景	The major pathway for the biosynthesis of phosphatidylcholine occurs via the CDP-choline pathway. The p protein encoded by this gene is the initial enzyme in the sequence and may play a regulatory role. The encoded protein also catalyzes the phosphorylation of ethanolamine. Two transcript variants encoding different isoforms have been found for this gene.
基因ID	1119

基因名	CHKA
Swiss	P35790
别名	CHKA;CHK;CK;CKI;EK

产品验证



Western blot analysis of CHKA expressed in Mouse brain,Rat testis,HepG2 using CHKA Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. B locking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn