

IL4R Rabbit pAb

货号: **AYP13952**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	25kDa/88kDa/89kDa
实测分子量	140kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,A-431,293T
细胞定位	Cell membrane,Secreted,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 26-232 of human IL4R (NP_000409.1).
序列	MKVLQEPTCVSDYMSISTCEWKMNGPTNCSTELRLLYQLVFLLEAHTCIPENGGAGCVCHLLMDDVVSADNYTLDLW AGQQLLWKGFSFKPSEHVKPRAPGNLTVHTNVSDTLTLLTWSNPYPPDNYLHLYAVNIWSENDPADFRIYNVTYLEPSLR IAASTLKSGISYRARVRAWAQCYNNTTWEWSPSTKWHNSYREPFEQH

靶点信息

研究背景	This gene encodes the alpha chain of the interleukin-4 receptor, a type I transmembrane protein that can bind interleukin 4 and interleukin 13 to regulate IgE production. The encoded protein also can bind interleukin 4 to promote differentiation of Th2 cells. A soluble form of the encoded protein can be produced by proteolysis of the membrane-bound protein, and this soluble form can inhibit IL4-mediated cell proliferation and IL5 upregulation by T-cells. Allelic variations in this gene have been associated with atopy, a condition that can manifest itself as allergic rhinitis, sinusitis, asthma, or eczema. Polymorphisms in this gene are also associated with resistance to human immunodeficiency virus type-1 infection. Alternate splicing results in multiple transcript variants.
基因ID	3566
基因名	IL4R
Swiss	P24394
别名	IL4R;CD124;IL-4RA;IL4RA

产品验证



Western blot analysis of IL4R expressed in K-562,A-431,293T using IL4R Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn