

ECHDC2 Rabbit pAb

货号: **AYP13955**

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	28kDa/31kDa
实测分子量	31kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse stomach,Mouse thymus,Mouse heart,Rat lung
细胞定位	Mitochondrion
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-292 of human ECHDC 2 (NP_001185890.1).
序列	MLRVLCLLRPWRPLRARGCASDGAAGGSEIQVRALAGPDQGITEILMNRPSARNALGNVVFSELLETLAQLREDRQVRVLLFRSGVKGVFCAGADLKEREQMSAEVGVFVQRLRGLMNDIAAFPAPTIAAMDGFALGGGLELALACDLRVAASSAVMGLIETTRGLLPGAGGTQRLPRCLGVALAKELIFTGRRLSGTEAHVLGLVNHAVAQNEEGDAAYQRARALAEILPQAPIAVRLGKVAIDRGTEVDIASGMAIEGMCYAQNIPTDRDRLEGMAAFREKRTPKFVGK

靶点信息

研究背景	Predicted to enable enoyl-CoA hydratase activity. Predicted to be involved in fatty acid beta-oxidation. Predicted to be active in mitochondrion.
基因ID	55268

基因名	ECHDC2
Swiss	Q86YB7
别名	ECHDC2

产品验证



Western blot analysis of ECHDC2 expressed in Mouse stomach,Rat lung,Mouse thymus using ECHDC2 R abbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn