

MED18 Rabbit pAb

货号: **AYP13957**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | WB: Mus musculus |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:1000 |
| 理论分子量 | 23kDa |
| 实测分子量 | 24KDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HT-29,Jurkat,Mouse liver,Mouse thymus |
| 细胞定位 | Nucleus |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-208 of human MED18 (NP_001120822.1). |
| 序列 | MEAPPVTMMPVTGGTINMMEYLLQGSVLDHSLESLIHRLRGLCDNMEPETFLDHEMVFLKGGQASPFLRARRSMDRA GAPWHLRYLGQPEMGDKNRHALVRNCVDIATSENLTDFLMEMGFRMDHEFVAKGHLFRKGIMKIMVYKIFRILVPGNTDS TEALSLSYLVESVVPAGQDMVSDDMKNFAEQLKPLVHLEKIDPKRLM |

靶点信息

| | |
|------|--|
| 研究背景 | MED18 is a component of the Mediator complex, which is a coactivator for DNA-binding factors that activate transcription via RNA polymerase II (Sato et al., 2003 [PubMed 12584197]).[supplied by OMIM, Oct 20 08] |
| 基因ID | 54797 |

| | |
|-------|-----------------|
| 基因名 | MED18 |
| Swiss | Q9BUE0 |
| 别名 | MED18;p28b;SRB5 |

产品验证



Western blot analysis of MED18 expressed in Mouse liver,HT-29,Jurkat using MED18 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blotting buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn