

# OAT3/SLC22A8 Rabbit pAb

货号: **AYP13959**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Mus musculus
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:100 - 1:500
理论分子量	11kDa/46kDa/49kDa/59kDa/60kDa
实测分子量	62KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney
细胞定位	Basolateral cell membrane,Multi-pass membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human OAT3/SLC22A8 (NP_004245.2).
序列	MTFSEILDRVGSMGHFQFLHVAIILGLPILNMANHLLQIFTAATPVHHCRRPPHNASTGPWVLPMPNGKPERCLRFVHPPN ASLPNDTQRAMEPCLDGWW

## 靶点信息

研究背景	This gene encodes a protein involved in the sodium-independent transport and excretion of organic anions, some of which are potentially toxic. The encoded protein is an integral membrane protein and appears to be localized to the basolateral membrane of the kidney. Multiple alternatively spliced transcript variants that encode different protein isoforms have been described for this gene.
基因ID	9376

基因名	SLC22A8
Swiss	Q8TCC7
别名	SLC22A8;OAT3

## 产品验证



Western blot analysis of OAT3/SLC22A8 expressed in Mouse kidney using OAT3/SLC22A8 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)