

# CCKBR Rabbit pAb

货号: **AYP13961**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Rattus norvegicus
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	41kDa/48kDa/55kDa
实测分子量	48kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,HT-29,K-562,BxPC-3,Mouse brain,Rat brain
细胞定位	Cell membrane,Multi-pass membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 240-335 of human CCKBR (NP_795344.1).
序列	LISRELYLGLRFDGSDSDSQSRVRNQGLPGAVHQNGRCRPETGAVGEDSDGCYVQLPRSRPALELTALTAPGPGSGS RPTQAKLLAKKRVRML

## 靶点信息

研究背景	This gene encodes a G-protein coupled receptor for gastrin and cholecystokinin (CCK), regulatory peptide s of the brain and gastrointestinal tract. This protein is a type B gastrin receptor, which has a high affinity for both sulfated and nonsulfated CCK analogs and is found principally in the central nervous system and the gastrointestinal tract. Alternative splicing results in multiple transcript variants. A misspliced transcrip t variant including an intron has been observed in cells from colorectal and pancreatic tumors.
基因ID	887

基因名	CCKBR
Swiss	P32239
别名	CCKBR;CCK-B;CCK2R;GASR

## 产品验证



Western blot analysis of CCKBR expressed in Mouse brain,Rat brain,A-549 using CCKBR Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane . Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)