

— ABLYBIO, Help Your Research



PHD2/EGLN1 Rabbit pAb

货号: **AYP13963**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	36kDa/43kDa/46kDa
实测分子量	50KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,A-549,Mouse heart
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human PHD2/EGLN1 (NP_071334.1).
------	---

靶点信息

研究背景	The protein encoded by this gene catalyzes the post-translational formation of 4-hydroxyproline in hypoxia-inducible factor (HIF) alpha proteins. HIF is a transcriptional complex that plays a central role in mammalian oxygen homeostasis. This protein functions as a cellular oxygen sensor, and under normal oxygen concentration, modification by prolyl hydroxylation is a key regulatory event that targets HIF subunits for proteasomal destruction via the von Hippel-Lindau ubiquitylation complex. Mutations in this gene are associated with erythrocytosis familial type 3 (ECYT3).
基因ID	54583
基因名	EGLN1
Swiss	Q9GZT9 (https://www.uniprot.org/uniprotkb/Q9GZT9/entry)
别名	EGLN1,C1orf12,ECYT3,HALAH,HIF-PH2,HIFPH2,HPH-2,HPH2,PHD2,SM20,ZMYND6,PHD2/EGLN1 Rabbit pAb,Hypoxia-inducible factor prolyl hydroxylase 2,Prolyl hydroxylase domain-containing protein 2,SM-20

产品验证



Western blot analysis of PHD2/EGLN1 expressed in 293T, A-549, Mouse heart using PHD2/EGLN1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)