

FBP2 Rabbit pAb

货号: **AYP13987**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	36kDa
实测分子量	36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse skeletal muscle,Rat Skeletal muscle,Rat kidney
细胞定位	Cell junction,Cytoplasm,Nucleus,Z line,myofibril,sarcomere
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-339 of human FBP2 (N P_003828.2).
序列	MTDRSPFETDMLTLTRYVMEKGRQAKGTGELTQLLNSMLTAIKAISSAVRKAGLAHLYGIAGSVNVTGDEVKKLDVLSNSL VINMVQSSYSTCVLVSEENKDAITAKEKRGKYVVCDFPLDGSSNIDCLASIGTIFAIYRKTSEDEPSEKDALQCGRNIVAAG YALYGSATLVALSTGQGVDLFMLDPALGEFVLVEKDVKIKKKGKIYSLNEG YAKYFDAATTEYVQKKKFPEDGSAPYGARY VGSMVADVHRTL VYGGIFLYPANQKSPKGKLRLLYECNPVAYIIIEQAGGLATTGTQPVLVDVKPEAIHQRVPLILGSPEDVQE YLTCVQKNQAGS

靶点信息

研究背景	This gene encodes a gluconeogenesis regulatory enzyme which catalyzes the hydrolysis of fructose 1,6-bi sphosphate to fructose 6-phosphate and inorganic phosphate.
基因ID	8789

基因名	FBP2
Swiss	O00757
别名	FBP2

产品验证



Western blot analysis of FBP2 expressed in Mouse skeletal muscle, Rat Skeletal muscle, Rat kidney using FBP2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn