— ABLYBIO, Help Your Research

Cyclin C Rabbit pAb

货号: AYP14017

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	**WB:** Mus musculus
应用	WB
推荐浓度	**WB:** 1:500 - 1:2000
理论分子量	22kDa/33kDa
实测分子量	33kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,NIH/3T3,Mouse liver,Mouse brain,Mouse lung,Mouse kidney,Rat liver
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-283 of human Cyclin C (NP_005181.2).

靶点信息

研究背景	The protein encoded by this gene is a member of the cyclin family of proteins. The encoded protein interacts with cyclin-dependent kinase 8 and induces the phophorylation of the carboxy-terminal domain of the large subunit of RNA polymerase II. The level of mRNAs for this gene peaks in the G1 phase of the cell cycle. Two transcript variants encoding different isoforms have been found for this gene.
基因ID	892
基因名	CCNC
Swiss	P24863 (https://www.uniprot.org/uniprotkb/P24863/entry)
别名	CCNC,CycC,SRB11,hSRB11,cyclin-C,Cyclin C Rabbit pAb,SRB11 homolog

产品验证



Western blot analysis of Cyclin C expressed in Mouse liver,Rat liver,MCF7 using Cyclin C Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)