

Cyclin C Rabbit pAb

货号: **AYP14017**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	22kDa/33kDa
实测分子量	33kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,NIH/3T3,Mouse liver,Mouse brain,Mouse lung,Mouse kidney,Rat liver
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-283 of human Cyclin C (NP_005181.2).
序列	MAGNFWQSSHYLQWILDKQDLLKERQKDLKFLSEEEYWKLIFFTNVIQALGEHLKLRQQVIATATVYFKRFYARYSLKSI DPVLMAPTCVFLASKVEEFGVVSNTRLIAAATSVLKTRFSYAFPKEFPYRMNHILECFYLLELMDCCCLIVYHPYRPLLQYVQD MGQEDMLLPLAWRIVNDTYRTDLCLLYPPFMIALACLHVACVVQKQDARQWFAELSDMEKILEIIRVILKLYEQWKNFDER KEMATILSKMPKPKPPPNSEGEQGPNGSQNSSYSQS

靶点信息

研究背景	The protein encoded by this gene is a member of the cyclin family of proteins. The encoded protein interacts with cyclin-dependent kinase 8 and induces the phosphorylation of the carboxy-terminal domain of the large subunit of RNA polymerase II. The level of mRNAs for this gene peaks in the G1 phase of the cell cycle. Two transcript variants encoding different isoforms have been found for this gene.
------	--

基因ID	892
基因名	CCNC
Swiss	P24863
别名	CCNC;CycC;SRB11;hSRB11;cyclin-C

产品验证



Western blot analysis of Cyclin C expressed in Mouse liver,Rat liver,MCF7 using Cyclin C Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn