

RBM14 Rabbit pAb

货号: **AYP14056**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | WB: Homo sapiens |
| 应用 | WB IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:1000 IF/ICC: 1:100 - 1:500 |
| 理论分子量 | 12kDa/13kDa/17kDa/37kDa/69kDa |
| 实测分子量 | 75KDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HeLa,293T,Jurkat,Rat testis |
| 细胞定位 | Nucleus,nucleolus |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 110-220 of human RBM 14 (NP_006319.1). |
| 序列 | VVKDYAFVHMEKEADAKAAIAQLNGKEVKGKRINVELSTKGQKKGPLAVQSGDKTKKPGAGDTAFPGTGGFSATFDYQ QAFGNSTGGFDGQARQPTPPFFGRDRSPLRRS |

靶点信息

| | |
|------|---|
| 研究背景 | This gene encodes a ribonucleoprotein that functions as a general nuclear coactivator, and an RNA splicing modulator. This protein contains two RNA recognition motifs (RRM) at the N-terminus, and multiple hexapeptide repeat domain at the C-terminus that interacts with thyroid hormone receptor-binding protein (TRBP), and is required for transcription activation. Alternatively spliced transcript variants encoding different isoforms (with opposing effects on transcription) have been described for this gene. |
|------|---|

| | |
|-------|------------------------------------|
| 基因ID | 10432 |
| 基因名 | |
| Swiss | Q96PK6 |
| 别名 | RBM14;COAA;PSP2;SIP;SYTIP1;TMEM137 |

产品验证



Western blot analysis of RBM14 expressed in Rat testis, HeLa, 293T using RBM14 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn