

UNC13D Rabbit pAb

货号: **AYP14058**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB
推荐浓度	WB: 1:1000 - 1:2000
理论分子量	41kDa/123kDa/128kDa
实测分子量	123kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,HeLa,HepG2,Mouse lung
细胞定位	Cytoplasm,Late endosome,Lysosome,Membrane,Peripheral membrane protein,Recycling endosome
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 891-1090 of human UNC13D (NP_954712.1).
序列	LIRKYFCSRIQQQAETTSEELGAVTVKASYRASEQKLRVELLSASSLLPLDSNGSSDPFVQLTLEPRHEFPELAARETQKHKKDLHPLFDETFEFLVPAEPCRKAGACLLLTVDYDTLGADDLEGEAFLPLREVPGLSGSEEPGEVPQTRLPLTYPAPNGDPILQLLEGRKGDREAQVFVRLRRHRAKQASQHALRPAP

靶点信息

研究背景	This gene encodes a protein that is a member of the UNC13 family, containing similar domain structure as other family members but lacking an N-terminal phorbol ester-binding C1 domain present in other Munc13 proteins. The protein appears to play a role in vesicle maturation during exocytosis and is involved in regulation of cytosolic granules secretion. Mutations in this gene are associated with familial hemophagocytic lymphohistiocytosis type 3, a genetically heterogeneous, rare autosomal recessive disorder.
------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

基因ID	201294
基因名	UNC13D
Swiss	Q70J99
别名	UNC13D;FHL3;HLH3;HPLH3;Munc13-4

产品验证



Western blot analysis of UNC13D expressed in Mouse lung,A-549,HeLa using UNC13D Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane . Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn