

— ABLYBIO, Help Your Research



# FCER1G Rabbit pAb

货号: **AYP14078**

## 产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Mus musculus <b>IHC:</b> Mus musculus
应用	WB
推荐浓度	<b>WB:</b> 1:1000 - 1:2000
理论分子量	9kDa
实测分子量	12kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse spleen
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 19-86 of human FCER1G (NP_004097.1).
------	--

## 靶点信息

研究背景	The high affinity IgE receptor is a key molecule involved in allergic reactions. It is a tetramer composed of 1 alpha, 1 beta, and 2 gamma chains. The gamma chains are also subunits of other Fc receptors.
基因ID	2207
基因名	FCER1G
Swiss	P30273 ( <a href="https://www.uniprot.org/uniprotkb/P30273/entry">https://www.uniprot.org/uniprotkb/P30273/entry</a> )
别名	FCER1G,FCRG,FCER1G Rabbit pAb,Fc receptor gamma-chain,Fc-epsilon RI-gamma,IgE Fc receptor subunit gamma

## 产品验证



Western blot analysis of FCER1G expressed in Mouse spleen using FCER1G Rabbit pAb at 1:1000. Second ary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)