

BST2 Rabbit pAb

货号: **AYP14120**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	18kDa/19kDa
实测分子量	35-40kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,HeLa,Mouse heart
细胞定位	Apical cell membrane,Cell membrane,Cytoplasm,GPI-anchor,Golgi apparatus,Late endosome,Lipid-anchor ,Membrane raft,Single-pass type II membrane protein,trans-Golgi network
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 49-161 of human BST2 (NP_004326.1).
序列	NSEACRDGLRAVMECRNVTHLLQQLTEAQKGFQDVEAQAATCNHTVMALMASLDAEKAQGQKKVEELEEITTLNHKL QDASAEVERLRRENQVLSVRIADKKYYPSSQDSS

靶点信息

研究背景	Bone marrow stromal cells are involved in the growth and development of B-cells. The specific function of the protein encoded by the bone marrow stromal cell antigen 2 is undetermined; however, this protein may play a role in pre-B-cell growth and in rheumatoid arthritis.
------	--

基因ID	684
基因名	BST2
Swiss	Q10589
别名	BST2;CD317;TETHERIN

产品验证



Western blot analysis of BST2 expressed in LO2, HeLa, Mouse heart using BST2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn