

PFDN2 Rabbit pAb

货号: **AYP14122**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IHC
推荐浓度	WB: 1:1000 - 1:5000 IHC: 1:100 - 1:200
理论分子量	16kDa
实测分子量	17kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	22Rv1,HeLa,293T,Jurkat,LO2,NIH/3T3
细胞定位	Cytoplasm,Mitochondrion,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-154 of human PFDN2 (NP_036526.2).
序列	MAENSGRAGKSSGSGAGKGA VSAEQVIAGFNRLRQEQRLASKAAELEMELNEHSLVIDTLKEVDETRKCYRMVGGVLV ERTVKEVLPALENNKEQIQKIETLTQQLQAKGKELNEFREKHNIRLMGEDEKPAAKENSEGAGAKASSAGVLVS

靶点信息

研究背景	This gene encodes a member of the prefoldin beta subunit family. The encoded protein is one of six subunits of prefoldin, a molecular chaperone complex that binds and stabilizes newly synthesized polypeptides, thereby allowing them to fold correctly. The complex, consisting of two alpha and four beta subunits, forms a double beta barrel assembly with six protruding coiled-coils.
基因ID	5202

基因名	PFDN2
Swiss	Q9UHV9
别名	PFDN2;PFD2

产品验证



Western blot analysis of PFDN2 expressed in 22Rv1, HeLa, 293T using PFDN2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn