

MXI1 Rabbit pAb

货号: **AYP14131**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	20kDa/21kDa/26kDa/32kDa
实测分子量	30kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,A-549,HepG2,Mouse brain,Rat heart,Rat brain
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 170-295 of human MXI1 (NP_569157.2).
序列	HTTLGLLNKAKAHIKKLEEAERKSQHLENLEREQFLKWRLEQLQGPQEMERIRMDSIGSTISSDRSDSEREEIEVDVEST EFSHGEVDNISTTSISDIDDHSSLPSIGSDEGYSSASVKLSFTS

靶点信息

研究背景	Expression of the c-myc gene, which produces an oncogenic transcription factor, is tightly regulated in normal cells but is frequently deregulated in human cancers. The protein encoded by this gene is a transcriptional repressor thought to negatively regulate MYC function, and is therefore a potential tumor suppressor. This protein inhibits the transcriptional activity of MYC by competing for MAX, another basic helix-loop-helix protein that binds to MYC and is required for its function. Defects in this gene are frequently found in patients with prostate tumors. Three alternatively spliced transcripts encoding different isoforms have been described. Additional alternatively spliced transcripts may exist but the products of these transcripts have not been verified experimentally.
基因ID	4601
基因名	MXI1
Swiss	P50539
别名	MXI1;MAD2;MXD2;MXI;bHLHc11

产品验证



Western blot analysis of MXI1 expressed in Mouse brain, Rat heart, U-87MG using MXI1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn