

— ABLYBIO, Help Your Research



NAGS Rabbit pAb

货号: **AYP14132**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IF: Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	58kDa
实测分子量	58kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,K-562,Mouse liver,Mouse small intestine,Mouse testis,Rat liver
细胞定位	Mitochondrion matrix
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 295-534 of human NAG S (NP_694551.1).
------	---

靶点信息

研究背景	The N-acetylglutamate synthase gene encodes a mitochondrial enzyme that catalyzes the formation of N-acetylglutamate (NAG) from glutamate and acetyl coenzyme-A. NAG is a cofactor of carbamyl phosphate synthetase I (CPSI), the first enzyme of the urea cycle in mammals. This gene may regulate ureagenesis by altering NAG availability and, thereby, CPSI activity. Deficiencies in N-acetylglutamate synthase have been associated with hyperammonemia.
基因ID	162417
基因名	NAGS
Swiss	Q8N159 (https://www.uniprot.org/uniprotkb/Q8N159/entry)
别名	NAGS,AGAS,ARGA,NAGS Rabbit pAb,Amino-acid acetyltransferase

产品验证



Western blot analysis of NAGS expressed in Mouse liver, Rat liver, 293T using NAGS Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)