

# SCRN2 Rabbit pAb

货号: **AYP14135**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Rattus norvegicus
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	41kDa/46kDa
实测分子量	47kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BxPC-3,HeLa,A375,Mouse liver,Mouse kidney,Mouse intestine
细胞定位	extracellular exosome
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 176-425 of human SCR N2 (NP_612364.2).
序列	GRLWAAQRIQEGARNISNQLSIGTDISAQHPELRTHAQAKGWWDGQGAFFDAQIFSLTQQPVRMEAAKARFQAGRELLR QRQGGITAEVMMGILRDKESGICMDSGGFRRTASMVSVLPQDPTQPCVHFLTATPDPSRSVFKPFIFGMGVAQAPQVLSP TFGAQDPVRTLPRFQTQVDRRHLYRGHQALGLMERDQDRGQQLQKQDLEQEGLEATQGLLAGEWAPPLWELGS LFQAFVKRESQAYA

## 靶点信息

研究背景	Predicted to enable cysteine-type exopeptidase activity and dipeptidase activity. Predicted to be involved in proteolysis. Located in extracellular exosome.
基因ID	90507

基因名	SCRN2
Swiss	Q96FV2
别名	SCRN2;Ses2

## 产品验证



Western blot analysis of SCRN2 expressed in Mouse liver, BxPC-3, HeLa using SCRN2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)