

— ABLYBIO, Help Your Research



NAPRT Rabbit pAb

货号: **AYP14173**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	56kDa/57kDa/60kDa
实测分子量	55kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,HT-29
细胞定位	Cytoplasm,cytosol
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 309-538 of human NAPRT1 (NP_660202.3).
------	--

靶点信息

研究背景	Nicotinic acid (NA; niacin) is converted by nicotinic acid phosphoribosyltransferase (NAPRT; EC 2.4.2.11) to NA mononucleotide (NaMN), which is then converted to NA adenine dinucleotide (NaAD), and finally to nicotinamide adenine dinucleotide (NAD), which serves as a coenzyme in cellular redox reactions and is an essential component of a variety of processes in cellular metabolism including response to stress (Hara et al., 2007).
基因ID	93100
基因名	NAPRT
Swiss	Q6XQN6 (https://www.uniprot.org/uniprotkb/Q6XQN6/entry)
别名	NAPRT,NAPRT1,PP3856,NAPRT Rabbit pAb,FHA-HIT-interacting protein,Nicotinate phosphoribosyltransferase domain-containing protein 1,FHIP

产品验证



Western blot analysis of NAPRT expressed in HepG2,HT-29 using NAPRT Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)