

— ABLYBIO, Help Your Research



MAD1/MAD1L1 Rabbit pAb

货号: **AYP14175**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	IHC: Mus musculus
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	72kDa/83kDa
实测分子量	95kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HepG2,MCF7
细胞定位	Chromosome,Cytoplasm,Nucleus,centromere,centrosome,cytoskeleton,kinetochore,microtubule organizing center,spindle
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 360-700 of human MAD1/MAD1/MAD1L1 (NP_001013859.1).
------	---

靶点信息

研究背景	MAD1L1 is a component of the mitotic spindle-assembly checkpoint that prevents the onset of anaphase until all chromosome are properly aligned at the metaphase plate. MAD1L1 functions as a homodimer and interacts with MAD2L1. MAD1L1 may play a role in cell cycle control and tumor suppression. Alternative splicing results in multiple transcript variants.
基因ID	8379
基因名	MAD1L1
Swiss	Q9Y6D9 (https://www.uniprot.org/uniprotkb/Q9Y6D9/entry)
别名	MAD1L1,MAD1,PIG9,TP53I9,TXBP181,MAD1/MAD1L1 Rabbit pAb,Mitotic arrest deficient 1-like protein 1,Mitotic checkpoint MAD1 protein homolog,Tax-binding protein 181

产品验证



Western blot analysis of MAD1/MAD1L1 expressed in HeLa,HepG2,MCF7 using MAD1/MAD1L1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)