

— ABLYBIO, Help Your Research



DAB1 Rabbit pAb

货号: **AYP14223**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IF/ICC
推荐浓度	WB: 1:100 - 1:500 IF/ICC: 1:50 - 1:200
理论分子量	24kDa/49kDa/57kDa/59kDa/63kDa
实测分子量	57KDa/59KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,Mouse small intestine,Rat brain
细胞定位	cytoplasm,cytosol,perinuclear region of cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 133-250 of human DAB 1 (NP_066566.3).
------	---

靶点信息

研究背景	The laminar organization of multiple neuronal types in the cerebral cortex is required for normal cognitive function. In mice, the disabled-1 gene plays a central role in brain development, directing the migration of cortical neurons past previously formed neurons to reach their proper layer. This gene is similar to disabled-1, and the protein encoded by this gene is thought to be a signal transducer that interacts with protein kinase pathways to regulate neuronal positioning in the developing brain.
基因ID	1600
基因名	DAB1
Swiss	O75553 (https://www.uniprot.org/uniprotkb/O75553/entry)
别名	DAB1,SCA37,DAB1 Rabbit pAb

产品验证



Western blot analysis of DAB1 expressed in U-87MG, Mouse small intestine, Rat brain using DAB1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)