

NR2F2 Rabbit pAb

货号: AYP14239

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IHC: Rattus norvegicus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	29kDa/31kDa/45kDa
实测分子量	47KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,Molt-4(Negative control),Mouse liver,PC-12,C6
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human NR2F2 (NP_066285.1).
序列	MAMVVSTWRDPQDEVPGSQGSQASQAPPVPGPPGAPHTPQTTPGQGGPASTPAQTAAGGQGGPGGPGSDKQQQQ HIECVCGDKSSGKHYGQFTCEGCK

靶点信息

研究背景	This gene encodes a member of the steroid thyroid hormone superfamily of nuclear receptors. The encoded protein is a ligand inducible transcription factor that is involved in the regulation of many different genes. Alternate splicing results in multiple transcript variants.
基因ID	7026

基因名	NR2F2
Swiss	P24468
别名	NR2F2;ARP1;CHTD4;COUPTFB;COUPTFII;NF-E3;SVP40;TFCOUP2

产品验证



Western blot analysis of NR2F2 expressed in Mouse liver, MCF7, Molt-4 (Negative control) using NR2F2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn