

# RBM22 Rabbit pAb

货号: **AYP14265**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>IP:</b> Homo sapiens
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	41kDa/46kDa
实测分子量	50KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,Mouse thymus,Rat lung
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-230 of human RBM22 (NP_060517.1).
序列	MATSLGSNTYNRQNWEDADFPILCQTCLGENPYIRMTKEYGKECKICARPFTVFRWCPCGVRMRFKKTEVCQTCSKLKNV CQTCLLDLEYGLPIQVRDAGLSFKDDMPKSDVNKEYYTQNMEREISNSDGTRPVGMLGKATSTSDMLLKLARTTPYYKRN RPHICSFVWKGECKRGEPCYRHEKPTDPPDPLADQNIKDRYYGINDPVADKLLKRASTMPRLDPPEDKT

## 靶点信息

研究背景	This gene encodes an RNA binding protein. The encoded protein may play a role in cell division and may be involved in pre-mRNA splicing. Related pseudogenes exist on chromosomes 6, 7, 9, 13, 16, 18, and X.
基因ID	55696

基因名	RBM22
Swiss	Q9NW64
别名	RBM22;Cwc2;ZC3H16;fSAP47

## 产品验证



Western blot analysis of RBM22 expressed in Raji, Mouse thymus, Rat lung using RBM22 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)